

HIV Trend Data: New diagnoses and risk factors



WASHTENAW COUNTY RESIDENTS

► Individuals newly diagnosed with HIV infection:

	2012	2013	2014	2015	2016	2017	2018
Total new HIV cases	24	33	28	27	27	15	29
▪ Male	23	29	26	26	23	15	25
▪ Female	1	4	2	1	4	0	4
▪ Black/African-American	8	18	10	15	11	3	14
▪ White	15	13	14	11	12	6	13
▪ Age 15 - 29 yr	14	24	18	19	12	9	12

Note: HIV+ data include anonymous and confidential cases, and exclude prison cases. Only cases that are newly diagnosed in Washtenaw County residents are included.

► Reported risk factors in individuals newly diagnosed with HIV:

	2012	2013	2014	2015	2016	2017	2018
MSM¹	88%	79%	89%	78%	59%	73%	83%
Bisexual male	29%	18%	21%	15%	15%	33%	7%
Injection drug use	0%	6%	4%	7%	7%	7%	0%
Hetero, high risk²	4%	3%	4%	4%	11%	0%	0%
Hetero, low risk³	8%	15%	11%	18%	22%	13%	14%


¹MSM = Men who have sex with men (includes bisexual males)

²Hetero, high risk = Heterosexual sex with IDU, known HIV+ partner, bisexual male, sex worker etc.

³Hetero, low risk = Heterosexual sex with no other risk identified

Note: Risk percentages are calculated in relation to the total number of cases. Risk percentages may add up to more or less than 100% due to some individuals having more than one risk and others having none identified.

► HIV burden in Blacks/African-Americans



Data source: CDC AtlasPlus

- Black/African-American men and women are living with HIV infection at a higher rate in Washtenaw County than White men or White women.
- In 2018, three of the new HIV diagnoses in Washtenaw were in Black/African-American women under the age of 30 years whose only risk factor was heterosexual contact.
- In 2018, all of the Black/African-American men diagnosed in Washtenaw reported having sex with men; several also reported sexual contact with women.